



Theoretical Competition, 5th IJSO, Gyeongnam, Korea, December 11, 2008

5th INTERNATIONAL JUNIOR SCIENCE OLYMPIAD

**THEORETICAL COMPETITION
December 11, 2008**

International Junior Science Olympiad

2008

7 ~ 16 December 2008

GYEONGNAM KOREA

Read carefully the following instructions:

1. The time available is 3 hours.
2. The total number of the problems is 3. Check that you have a complete set of the test problems and the answer sheets.
3. Use only the pen provided.
4. Write down your name, code, country, and signature in the first page of your answer sheet. Write down your name and code in the other pages of your answer sheet.
5. Read carefully each problem and write the correct answer in the answer sheet.
6. All competitors are not allowed to bring any stationary and tools provided from outside. After completing your answers, all of the question and answer sheets should be put neatly on your desk.
7. Point rules : According with each question marking.


EXAMINATION RULES

1. All competitors must be present at the front of examination room ten minutes before the examination starts.
2. No competitors are allowed to bring any tools except his/her personal medicine or any personal medical equipment.
3. Each competitor has to sit according to his or her designated desk.
4. Before the examination starts, each competitor has to check the stationary and any tools (pen, ruler, calculator) provided by the organizer.
5. Each competitor has to check the question and answer sheets. Raise your hand, if you find any missing sheets. Start after the bell.
6. During the examination, competitors are not allowed to leave the examination room except for emergency case and for that the examination supervisor will accompany them.
7. The competitors are not allowed to bother other competitor and disturb the examination. In case any assistance is needed, a competitor may raise his/her hand and the nearest supervisor will come to help.
8. There will be no question or discussion about the examination problems. The competitor must stay at their desk until the time allocated for the examination is over, although he/she has finished the examination earlier or does not want to continue working.
9. At the end of the examination time there will be a signal (the ringing of a bell). You are not allowed to write anything on the answer sheet, after the allocated time is over. All competitors must leave the room quietly. The question and answer sheets must be put neatly on your desk.

Problem I

Pressure

In 1643, Torricelli filled a 1 m long tube, sealed at one end, with mercury and inverted it into a basin containing mercury. The column of mercury fell to about 76 cm above the level of the mercury basin, leaving a ‘Torricellian’ vacuum above. After this discovery, it has been widely accepted that 1 atm (atmospheric pressure) is equivalent to 760 mmHg or 760 Torr. Pressure is defined as ‘the force per unit area applied to an object in a direction perpendicular to the surface’. Assume that the density of mercury is 13.6 times density of water ($1.0 \times 10^3 \text{ kg/m}^3$) and g is 9.8 m/s^2 .



I-1) Express the unit of the pressure in terms of kg, m, and s. (0.3 point)

I-2) How high does a water column go up when you use a sufficiently long tube filled with water instead of mercury at 1 atm? (0.5 point)

I-3) What is the value of 1 atm in SI units? (0.5 point)

Blood Pressure

Blood pressure refers to the force per unit area exerted by circulating blood on the walls of blood vessels, and constitutes one of the principal vital signs. The term blood pressure generally refers to arterial pressure, i.e., the pressure in the larger arteries (the blood vessels that take blood away from the heart). During the heart pumping cycle arterial pressure reaches a maximum and a minimum value, and these pressures are called systolic and diastolic and for a healthy heart are around 120 mmHg (systolic) and 80 mmHg (diastolic) measured at heart level.

I-4) If a pilot, who has a healthy heart, is accelerating upward (in the direction of his/her head), estimate the minimum acceleration for which blood supply to the pilot’s brain stops

totally?

(Assume the followings: 1. The blood pressure does not change. 2. The density of blood is the same as that of water. 3. The brain is located 42 cm above the heart. 4. The air pressure in the cockpit remains constant.) (1.2 points)

I-5) The heart pumps blood into the aorta (main artery), which has an inner radius of 1.2 cm. The aorta feeds 32 major arteries. If blood in the aorta travels at a speed of 25 cm/s, **at approximately what speed does it travel in the 32 major arteries?** (Assume that blood can be treated as an incompressible and non-viscous fluid and that each artery has an inner radius of 0.2 cm.) (1.0 point)

Poiseuille's law

In fact, blood is a viscous fluid. The volume flow rate $\Delta V/\Delta t$ for streamline flow of a viscous fluid through a horizontal, cylindrical pipe is known to be

$$\Delta V/\Delta t \propto \Delta P r^4,$$

where r and ΔP are the inner radius of the pipe and the pressure difference between the ends of the pipe.

I-6) A cardiologist reported to his patient that the radius of one of the major arteries decreased by 10.0% as compared to a normal condition. **What percent increase in the blood pressure drop across the artery is required to maintain the normal blood flow through this artery?** Assume that the blood flow is streamline flow. (1.0 point)

Analogy to electric circuits

The blood circulatory system is similar to an electric circuit in certain ways; below is a table of the corresponding elements in the analogy between the blood circulatory system and an electric circuit.

I-7) Choose a number among 1 to 5 to fit as correct analogy in each blank (A) to (E). (0.2 point each)

blood circulating system	electric circuit
heart	(A)
blood	(B)
blood pressure	(C)
blood vessel	(D)
blood flow	(E)

1	charge
2	electric potential
3	wire
4	battery
5	electric current


Kirchhoff's rules of circuits

By this analogy, we can understand Kirchhoff's rules of circuits.

1. The sum of the currents that flow into a junction is equal to the sum of the currents that flow out of the same junction.
2. If a closed path is followed in a circuit, the algebraic sum of the potential changes must be zero.

Using Kirchhoff's rules, solve the following problems.

The Wheatstone bridge is a circuit used to measure unknown resistances. The bridge in the figure is balanced: no current flows through the galvanometer (or ammeter, **G**). (Assume the galvanometer is ideal: internal resistance is negligible compared to other resistors)



I-8) What is the potential difference (voltage) between the points A and B? (0.5 point)

I-9) What is the resistance of R_x ? (0.5 point)

If R_x is $6\ \Omega$, the current through the galvanometer is 0.2 A. The current flows A to B.

I-10) What is the voltage between the points A and B? (0.5 point)

I-11) What is the voltage of the battery? (1.5 points)

I-12) If you disconnect the galvanometer, what is the voltage between the points A and B? And which point is higher? (1.5 points)

Problem II Various energy sources

Gasoline is the most commonly used mixture of hydrocarbon compounds as a fuel for motor vehicles. The major component of gasoline is octane.

Liquefied petroleum gas (LPG) is a mixture of low molecular weight hydrocarbons condensed into liquid under high pressure. LPG is a mixture of propane and butane in different ratios.

Liquefied natural gas (LNG) is natural gas mainly consisting of methane.

These compounds can be used as energy sources since they release large amount of heat by combustion reactions with oxygen (O_2). The following table provides useful data about these fuels. (Assume that all gases are ideal gases and the volume of the liquid does not change with temperature.)

Fuel	Major component	Chemical formula	Density of liquid (kg/L)	Energy content (kJ/kg)
Gasoline	Octane	C_8H_{18}	0.70	44000
LPG (liquefied petroleum gas)	Propane	C_3H_8	0.50	46000
	Butane	C_4H_{10}	0.57	46000
LNG (liquefied natural gas)	Methane	CH_4	0.42	54000

II-1) In some countries, the engine efficiency is represented by the amount of fuel consumed for the vehicle to travel 100 km, while in other countries by the distance for the vehicle to travel with 1.0 L of fuel. A car has a gasoline consumption of 13.0 L/100 km. If LPG can be used as an alternative fuel for this car, **calculate the distance that the car can travel with 1.0 L of liquid LPG.** Assume that LPG consists of pure propane, and the engine efficiency of the car is the same for both gasoline and LPG. (2.5 points)

II-2) LNG is usually stored as a liquid at very low temperature. If 10 mL of liquid LNG is completely vaporized to a gaseous state at 25° and 1.0 atm, **what is the volume of the resulting gaseous LNG?** Assume that LNG is pure methane. (1.0 point)

II-3) LPG is commonly prepared as a mixture of propane and butane in different proportions. A cylinder of LPG contains both propane and butane in the mass ratio of 3:2. **What will be the density of gas mixture when LPG in this cylinder is completely vaporized to a gaseous state at 25 °C and 1.0 atm? (1.5 points)**

II-4) Hydrocarbon fuels produce carbon dioxide (CO₂) and water vapor when they undergo complete combustion. **Write the balanced chemical equation for the complete combustion of octane. (1 point)**

II-5) CO₂ gas produced during combustion of fuels is one of the major greenhouse gases. If the 1.0 kJ of energy is obtained by burning octane or methane, **what is the mass of CO₂ produced from each fuel?** Assume that both fuels undergo complete combustion. (2.5 points)

II-6) Because fossil fuels are limited resources and cause environmental problems, a great deal of efforts has been made to utilize solar energy as a new clean energy source. The solar cell is a device converting the light energy of the sun into electric energy. We want to obtain the same amount of energy from the solar cell in **1.0 hour** as that produced from combustion of 1.0 L of liquid gasoline. Assume that 1000 W solar power reaches 1.0 m² area of Earth surface. If the conversion efficiency of the solar cell is 20 %, **what is the minimum surface area of the solar cell?** (1.5 points)

Reference data :

Atomic mass: H = 1, C = 12, N = 14, O = 16 amu

Gas constant $R = 0.082 \text{ L atm mol}^{-1} \text{ K}^{-1} = 8.3 \text{ J mol}^{-1} \text{ K}^{-1}$

The molar gas volume at 25 °C (room temperature) and pressure (1.0 atm) is 24.5 L/mol (24.5 dm³/mol).

Problem III

Photoperiodism and Control of Flowering


The environmental stimulus that plants use most often to detect the time of year is the photoperiod, the relative lengths of night and day in a 24-hour daily cycle. Plants whose flowering is triggered by photoperiod fall into two groups. One group, the short-day plants, generally flowers during late summer, autumn, or winter, when light periods are significantly shorter. In contrast, flowers of long-day plants usually bloom in late spring or early summer, when light periods are relatively longer. In the 1940s, researchers have discovered that photoperiodic response of flowering is actually controlled by the length of night (length of darkness), not the length of day (length of light). In fact, the so-called ‘short-day’ plants are actually ‘long-night’ plants.

To flower, short-day plants require a certain night length which is longer than a critical length to flower. For instance, the common tobacco (*Nicotiana tabacum*), which is a typical short-day plant, will flower when the night length is at least a critical length of darkness or longer. Long-day plants only flower when the night length is shorter than their critical night length. Henbane (*Hyoscyamus niger*), which is an example of a long-day plant, will flower when night length is shorter than 14 hours. Plants detect the night length very precisely; some short-day plants will not flower if night length is even 10 minutes shorter than the necessary critical length. Some plant species always flower on the same day each year.

Flowers of some plants bloom after a single exposure to the photoperiod required for flowering. Researchers found that if the night-time part of the photoperiod is interrupted by even a few minutes of dim light, short-day plant will not flower. Chrysanthemums are short-day plants that normally flower in the autumn, but their blooming can be stalled until the following spring by interrupting each long night with a flash of red light, thus, as a result, turning one long night into two short nights.


III-1~3. (1.0 point each)

Figure 1.



III-1) When a flash of light is given during the night portion of the photoperiod as in Figure 1, does henbane flower?

III-2) When the following condition is given, does tobacco flower?




III-3) When the following condition is given, does henbane flower?



III-4~5. (1.0 point each)

How does a plant actually measure photoperiod? Pigments called phytochromes are part of the answer. Phytochrome is a photoreceptor that specifically detects red light. Researchers learned that red light is the most effective light for interrupting the night length. Phytochrome is photoreversible, reverting back and forth between two isomers, depending on the color of light provided. The P_r isomer form of phytochrome shows maximum absorption of red light (wavelength, 660 nm), whereas the P_{fr} isomer mainly absorbs far-red light (wavelength, 730 nm). This interconversion between two molecular isomers is a switching mechanism that controls various light-induced events in the life of the plant, including flowering in short-day and long-day plants.

Figure 2.



If a flash of red light (R) during darkness is followed by a flash of far-red light (F), short-day plants will flower as in fig. 2c in Figure 2.

III-4) If a flash of red light (R) during darkness is followed by a flash of far-red light (F), do you expect the long-day plant to flower in the same conditions as in fig. 2c?

The sequential R-F-R light treatment results in the same effect as in a single R treatment. Thus, short-day plant will not flower as in fig. 2d.

III-5) When the same sequential R-F-R light treatments are given to the long-day plant, do you expect the long-day plant to flower in the same condition as in Figure 2-d?

III-6, 7. Fill in the following blanks with either P_R or P_{fr} . (1.0 point)

Each day, the conversion of the P_{fr} to the P_R isoform mainly occurs during the continuous darkness that follows sunset, even if far-red light is not present. After sunrise, however, much of the phytochrome is rapidly converted from the P_R isoform to the P_{fr} isoform because sunlight is significantly richer in red light than far-red light.


Phytochrome helps the plant distinguish daytime and night-time because phytochromes are mainly in the (III-6. _____) isoform during the day and mainly in the (III-7. _____) isoform during the night.

III-8~9 (1.5 points each)


Although flowers are formed from apical buds, it is the plant's leaves that detect changes in photoperiod and produce signaling molecules that cue buds to develop as flowers. It is reported that the signaling molecules are able to move through the stem. When grown individually under short-day conditions, a short-day plant will flower whereas a long-day plant will not. In the following classic experiment, a long-day plant is grafted to the short-day plant as in Figure 3.

Answer the following questions.

Figure 3.



III-8) Do you expect the grafted plant to flower under the condition shown below?



III-9) Do you expect the grafted plant to flower under the condition shown below?

